ANEURYSMAL DILATATION OF THE RIGHT ATRIUM (RA)

Prepared by **Dr. Sidhartha Mullapudi**, DrNB (Cardiology), MD (General Medicine)
Last reviewed: October 2026

I ETIOLOGY AND CLASSIFICATION (CAUSES)

- **Core Principle:** Aneurysmal dilatation of the right atrium—also termed **Idiopathic Dilatation of the Right Atrium (IDRA)**, congenital right atrial aneurysm, or Giant Right Atrium—is a rare congenital anomaly characterized by massive enlargement of the right atrium in the absence of primary right ventricular myocardial disease, pulmonary hypertension, or intrinsic tricuspid valve disease. When underlying hemodynamic or structural pathology is present, it is classified as secondary dilatation.
- **Anatomical Subtypes:** Primary congenital malformations of the right atrium include diffuse global enlargement (**IDRA**), focal saccular **Right Atrial Diverticulum** (single or multiple outpouchings of the RA free wall), and isolated **Right Atrial Appendage Aneurysm (RAAA)**.

A Primary (Idiopathic / Congenital RA Aneurysm)

- **Definition:** A congenital structural anomaly in which the right atrial wall is focally or diffusely thinned and dilated due to an intrinsic defect of the atrial myocardium (histologically showing focal absence or dysplasia of the pectinate muscles, lipomatous degeneration, or endocardial fibrosis).
- **Diagnostic Rule of Exclusion:** IDRA is strictly a diagnosis of exclusion. Primary tricuspid leaflet anatomy, right ventricular myocardium, and pulmonary arterial pressures must be normal. However, mild-to-moderate **secondary (functional) tricuspid regurgitation** may develop over time due to progressive tricuspid annular dilation from the giant atrium.

B Secondary (Hemodynamic / Acquired Dilatation)

- **Volume Overload Lesions:** Severe primary **Tricuspid Regurgitation** (classically **Ebstein Anomaly**, where the atrialized right ventricle and functional RA combine to produce massive right atrial enlargement), large **Atrial Septal Defects (ASD)**, **Partial or Total Anomalous Pulmonary Venous Connection (PAPVC/TAPVC)**, or **Gerbode Defect** (LV-to-RA shunt).
- **Pressure Overload Lesions:** Severe **Pulmonary Hypertension**, severe **Pulmonary Valve Stenosis**, tricuspid stenosis, or chronic cor pulmonale.
- **Myocardial and Restrictive Diseases: Arrhythmogenic Right Ventricular Cardiomyopathy (ARVC)**, right-sided **Restrictive Cardiomyopathies** (classically **Right Ventricular Endomyocardial Fibrosis [RV-EMF]**, which produces a giant right atrium with fibrous obliteration of the RV apex, or cardiac amyloidosis), and **Uhl Anomaly** (congenital parchment-like absence of RV myocardium).

II CLINICAL PRESENTATION AND COMPLICATIONS

- **Incidental Presentation:** Approximately 45 to 50 percent of patients with IDRA are completely asymptomatic and diagnosed incidentally on fetal echocardiography or routine postnatal chest radiography showing marked cardiomegaly.
- **The Classic Complication Triad:**
  - **Supraventricular Arrhythmias (Most Common):** Progressive stretching and patchy fibrosis of the atrial wall disrupt electrical conduction, triggering **Atrial Flutter**, **Atrial Fibrillation**, focal atrial tachycardia, or atrioventricular reciprocating tachycardia via associated accessory pathways (rarely causing sudden cardiac death).
  - **Thromboembolism:** Severe blood stasis within the giant atrial cavity predisposes to right atrial mural thrombus formation, causing recurrent **Pulmonary Embolism (PE)** or paradoxical systemic embolism across a stretched patent foramen ovale.
  - **Mechanical Compression Syndromes:** A massively dilated right atrium can compress the **Right Ventricle** and inflow tract (impairing RV filling and mimicking cardiac tamponade or right heart failure) or compress adjacent lung parenchyma and bronchi (causing dyspnea or airway compression/atelectasis in infants).

III DIAGNOSTIC EVALUATION

- **Electrocardiogram (ECG):** Demonstrates tall, peaked P waves (**P-pulmonale** greater than 2.5 mm in inferior leads II, III, and aVF) when in sinus rhythm, incomplete right bundle branch block, or atrial tachyarrhythmias (atrial flutter or fibrillation), notably without voltage criteria for right ventricular hypertrophy in pure IDRA.
- **Chest X-Ray (CXR):** Shows dramatic cardiomegaly (cardiothoracic ratio often exceeding 0.65 to 0.80) with a massively prominent, convex right heart border and normal pulmonary vascular markings, frequently misdiagnosed initially as a large pericardial effusion or Ebstein anomaly.
- **Echocardiography (TTE and TEE):** First-line imaging modality demonstrating a massively dilated right atrium. TTE confirms normal apical insertion of the septal tricuspid leaflet (displacement less than 8 mm/m2 body surface area, excluding Ebstein anomaly) and normal RV systolic function. **Transesophageal Echocardiography (TEE)** is highly sensitive for ruling out a sinus venosus or secundum ASD, anomalous pulmonary venous drainage, and detecting **spontaneous echo contrast (“smoke”)** or mural thrombus.
- **Cardiac MRI (CMR):** Reference standard for volumetric quantification and myocardial tissue characterization. CMR rules out RV fibrofatty replacement and Late Gadolinium Enhancement (**ARVC**), RV apical obliteration (**Endomyocardial Fibrosis**), or complete absence of RV myocardium (**Uhl anomaly**), and definitively distinguishes a thin-walled RA aneurysm from a loculated pericardial effusion, pericardial cyst, or anterior mediastinal tumor.

IV MANAGEMENT STRATEGY

A Conservative / Medical Management

- **Target Patient:** Asymptomatic patients with incidental, non-progressive mild-to-moderate enlargement and no history of arrhythmias, intracavitary thrombus, or compressive symptoms.
- **Surveillance:** Close clinical, ECG, and echocardiographic follow-up every 6 to 12 months, alongside antiarrhythmic drug therapy for controllable supraventricular arrhythmias.
- **Anticoagulation:** Oral anticoagulation (**Warfarin** or **DOACs**) is indicated in patients with atrial fibrillation or flutter, documented mural thrombus, prior pulmonary embolism, or severe right atrial stasis (spontaneous echo contrast) even in sinus rhythm to prevent pulmonary thromboembolism (low-dose aspirin is sometimes used empirically in asymptomatic children without stasis).

B Surgical Management (Right Atrial Reduction Atrioplasty)

- **Target Patient:** Symptomatic patients with recurrent or refractory atrial arrhythmias, intracavitary thrombus or pulmonary embolism despite anticoagulation, mechanical compression of the right ventricle or airway, severe functional tricuspid regurgitation, or rapid aneurysmal expansion.
- **Surgical Components:**
  - **Reduction Atrioplasty:** Surgical excision of the redundant, aneurysmal, paper-thin right atrial wall on cardiopulmonary bypass to restore physiological RA geometry and volume.
  - **Concomitant Maze Procedure:** In patients with atrial fibrillation or flutter, a right-sided **Cryo-Maze or Radiofrequency Maze procedure** is performed simultaneously to interrupt macro-reentrant arrhythmia circuits.
  - **Concomitant Valve Repair:** Tricuspid ring annuloplasty is performed if annular dilation has caused significant functional tricuspid regurgitation; in secondary dilatation due to Ebstein anomaly, tricuspid valve repair (**Cone reconstruction**) or replacement is performed simultaneously with RA reduction.

V SUMMARY TABLE FOR EXAM

Disease State	Distinguishing Feature	Primary Management
Primary RA Aneurysm (IDRA)	Normal septal TV insertion (less than 8 mm/m2); normal PA pressure and RV myocardium	Anticoagulation if stasis/AF; surgical **Reduction Atrioplasty + Maze** if symptomatic
Ebstein Anomaly	Apically displaced septal/posterior TV leaflets (8 mm/m2 or greater); atrialized RV	**Cone tricuspid repair** (or replacement) + RA reduction plasty
Endomyocardial Fibrosis (RV-EMF)	Restrictive RV physiology; fibrous obliteration of RV apex with giant RA	Diuretics and anticoagulation; surgical endocardiectomy + TV repair / Glenn shunt
ARVC / Uhl Anomaly	RV akinesia/aneurysms and LGE on CMR (ARVC); parchment-thin RV wall (Uhl)	Antiarrhythmics/ablation and **ICD** in ARVC; avoid isolated RA plasty
Severe Pulmonary Hypertension	Elevated PA pressures on RHC; RV hypertrophy and D-shaped LV	Treat underlying **PAH**; isolated RA reduction plasty is contraindicated

KEY REFERENCES

- **Binder TM, Rosenhek R, Frank H, Gwechenberger M, Maurer G, Baumgartner H:** Congenital malformations of the right atrium and the coronary sinus: an analysis based on 103 cases reported in the literature and four additional cases, *Chest*, 2000.
- **Baumgartner H, De Backer J, Babu-Narayan SV, et al.:** 2020 ESC Guidelines for the management of adult congenital heart disease, *European Heart Journal*, 2020.
- **Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD:** *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine*, 12th Ed., Elsevier, 2022.