CORONARY COLLATERAL CIRCULATION

Prepared by **Dr. Sidhartha Mullapudi**, DrNB (Cardiology), MD (General Medicine)
Last reviewed: October 2026

I CORE CONCEPTS AND ANATOMY

- **Definition:** Coronary collaterals are endogenous inter-arterial anastomotic channels ("natural bypasses") that connect different segments of the same epicardial coronary artery (intracoronary or homocoronary collaterals) or connect branches of one epicardial coronary artery to another (intercoronary collaterals, such as septal connections between the LAD and PDA, or atrial/epicardial connections between the LCx and RCA).
- **The Pre-Existing "Dormant" Network:** Unlike skeletal muscle, the human heart possesses an innate network of pre-existing microscopic anastomotic channels (40 to 200 micrometers in diameter) at birth. Under normal physiological conditions without coronary stenosis, there is no trans-anastomotic pressure gradient across these vessels, so net blood flow is negligible and they remain angiographically invisible (as standard coronary angiography only resolves vessels greater than 100 micrometers).
- **The Recruitment Trigger:** When a severe epicardial stenosis (typically greater than 70 to 80 percent luminal diameter narrowing) or a Chronic Total Occlusion (CTO) develops, distal post-stenotic coronary perfusion pressure falls, creating a steep pressure gradient between the donor artery and the recipient territory that recruits and remodels these dormant conduits into functional collateral arteries (up to 100 to 800 micrometers in diameter).

II PATHOPHYSIOLOGY OF COLLATERAL GROWTH (ANGIOGENESIS VS. ARTERIOGENESIS)

- **Angiogenesis (Hypoxia-Driven Capillary Sprouting):** The sprouting of new thin-walled endothelial capillary networks from pre-existing post-capillary venules. Driven directly by myocardial ischemia and tissue hypoxia, which stabilize Hypoxia-Inducible Factor-1 alpha (HIF-1a) and upregulate Vascular Endothelial Growth Factor (VEGF) and Fibroblast Growth Factor (FGF). While angiogenesis increases local capillary density, capillaries lack smooth muscle media and cannot conduct high bulk flow to replace an occluded epicardial artery.
- **Arteriogenesis (Shear Stress-Driven Conduit Remodeling):** The dominant mechanism of functional human coronary collateral development. Driven primarily by a biomechanical force—increased fluid shear stress across pre-existing arteriolar anastomoses due to the trans-stenotic pressure gradient (rather than local hypoxia alone).
- **Cellular Cascade:** Elevated endothelial shear stress upregulates endothelial Nitric Oxide Synthase (eNOS), Monocyte Chemoattractant Protein-1 (MCP-1), and adhesion molecules (ICAM-1/VCAM-1), recruiting circulating monocytes that differentiate into perivascular macrophages, secrete Matrix Metalloproteinases (MMPs) and growth factors (GM-CSF, TGF-beta, FGF-2), and induce vascular smooth muscle cell proliferation—enlarging the vessel diameter up to 10- to 20-fold into a three-layered muscular artery.

III ANGIOGRAPHIC AND PHYSIOLOGICAL GRADING OF COLLATERALS

A The Rentrop Classification (Angiographic Filling of Recipient Vessel)

- **Grade 0:** No visible angiographic filling of any collateral channels.
- **Grade 1:** Filling of side branches of the occluded/recipient artery via collateral channels, without visualization of the main epicardial segment.
- **Grade 2:** Partial filling of the main epicardial trunk of the occluded/recipient artery via collateral channels.
- **Grade 3:** Complete filling of the main epicardial trunk of the occluded/recipient artery via collateral channels.

B The Werner Classification (Collateral Connection [CC] Size — Standard for Retrograde CTO-PCI)

- **CC0:** No continuous connection between donor and recipient artery.
- **CC1:** Continuous, thread-like connection (diameter approx. 0.3 mm or less; most common channel crossed in retrograde CTO-PCI).
- **CC2:** Continuous, small side-branch-like connection throughout its course (diameter approx. 0.4 mm or greater).

C Physiological Quantitative Index: Collateral Flow Index (CFI)

- **Formula and Cut-off:** The invasive gold standard for quantifying collateral function during a 1-minute balloon occlusion: Pressure-derived CFI = (Pw - CVP) / (Pa - CVP), where Pw is distal coronary wedge pressure during balloon occlusion, Pa is mean aortic pressure, and CVP is central venous pressure. A CFI of 0.25 or greater indicates well-developed collaterals sufficient to prevent ECG signs of myocardial ischemia during acute coronary occlusion (present in roughly 20 to 25 percent of patients with CAD).

Classification System	Parameter Assessed	Grades & Key Clinical Utility
Rentrop Classification (1985)	Extent of contrast opacification of the recipient epicardial artery	Grades 0 to 3; predicts myocardial viability and infarct size limitation
Werner Classification (2003)	Anatomical size and continuity of the collateral channel itself	CC0, CC1 (thread-like), CC2 (branch-like); guides wire/microcatheter tracking in Retrograde CTO-PCI
Collateral Flow Index (CFI)	Hemodynamic pressure/velocity ratio during balloon occlusion	CFI = (Pw - CVP) / (Pa - CVP); CFI 0.25 or greater prevents acute ischemic ST-elevation

IV CLINICAL SIGNIFICANCE AND HEMODYNAMIC EFFECTS

- **Myocardial Viability and Infarct Size Limitation:** In acute STEMI or chronic total occlusion, robust pre-existing collaterals (Rentrop Grade 2 or 3 or CFI 0.25 or greater) maintain resting myocardial perfusion (approx. 20 to 30 percent of maximal flow), limiting transmural infarct size, preserving hibernating myocardium, reducing no-reflow, preventing left ventricular aneurysm formation, and lowering cardiogenic shock and mortality rates.
- **The "Coronary Steal" Phenomenon:** Even well-developed Rentrop Grade 3 collaterals can rarely increase coronary flow more than 2-fold during exercise or pharmacological vasodilation (adenosine/dipyridamole). During hyperemia or exercise, arteriolar resistance in the normal donor territory drops sharply while microvascular resistance in the collateral-dependent recipient bed is already maximally vasodilated at rest; blood is shunted away from the collateral pathway into the donor bed, causing a drop in distal collateral perfusion pressure (coronary steal) and effort angina.

V THE PARADIGM SHIFT IN CTO REVASCULARIZATION (ESC 2024 CCS GUIDELINES)

- **The Physiological Reality:** Angiographic Rentrop Grade 3 collaterals only demonstrate that contrast reaches the distal vessel at rest; invasive FFR measurements show that over 90 percent of CTO territories supplied by Rentrop Grade 3 collaterals remain physiologically ischemic (FFR less than 0.80) during demand.
- **Current Evidence and Guideline Stance (EUROCTO, IMPACTOR-CTO, DECISION-CTO, REVASC, EXPLORER):** Randomized trials (EUROCTO, IMPACTOR-CTO) demonstrate that CTO-PCI improves angina relief, ischemic burden, and quality of life (Seattle Angina Questionnaire) compared with optimal medical therapy (OMT) alone, whereas DECISION-CTO showed no reduction in hard clinical events, REVASC showed no improvement in regional LV function, and EXPLORER (CTO-PCI of a non-infarct artery after STEMI) showed no improvement in LVEF. Consequently, the 2024 ESC Chronic Coronary Syndromes Guidelines recommend CTO-PCI (Class IIa) for patients with refractory angina or a large area of documented ischemia/viability despite optimal antianginal therapy, rather than for asymptomatic patients based on anatomy alone.

VI PRACTICAL CONSIDERATIONS IN THE INDIAN CLINICAL CONTEXT

- **Impaired Arteriogenesis in Indian Diabetics:** Indian patients have a high prevalence of premature Diabetes Mellitus and insulin resistance, which blunts shear-stress-mediated eNOS activation, induces VEGF resistance in monocytes (the Waltenberger defect), and increases endothelial dysfunction—resulting in poorer collateral recruitment, larger infarct sizes, and diffuse multivessel disease.
- **Retrograde CTO-PCI Conduits:** In high-volume Indian interventional centers, septal perforator collaterals (safer, intramyocardial; perforation causes contained septal hematoma rather than tamponade) and epicardial collaterals (higher perforation/tamponade risk) are graded using the Werner CC classification for retrograde strategy planning (collateral grade is not part of the J-CTO Score) to perform retrograde wire crossing and Reverse CART (Controlled Antegrade and Retrograde Tracking).
- **Role of EECP in "No-Option" Refractory Angina:** Enhanced External Counterpulsation (EECP)—which inflates sequential lower-extremity pneumatic cuffs during diastole to augment aortic diastolic pressure and coronary shear stress—is utilized (Class IIb) as a non-invasive adjunct for symptom relief in patients with diffuse, non-revascularizable CAD ("no-option" refractory angina).

KEY REFERENCES

- **Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD:** *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine*, 12th Ed., Elsevier, 2022.
- **Rentrop KP, Cohen M, Blanke H, Phillips RA:** Changes in collateral channel filling immediately after controlled coronary artery occlusion by an angioplasty balloon in human subjects, *Journal of the American College of Cardiology*, 1985.
- **Werner GS, Ferrari M, Heinke S, et al.:** Angiographic assessment of collateral connections in comparison with invasively determined collateral function in chronic coronary occlusions, *Circulation*, 2003.
- **Vrints C, Andreotti F, Koskinas KC, et al.:** 2024 ESC Guidelines for the management of chronic coronary syndromes, *European Heart Journal*, 2024.