INDICATIONS OF CARDIAC CT IN CONGENITAL HEART DISEASE (CHD)

Prepared by **Dr. Sidhartha Mullapudi**, DrNB (Cardiology), MD (General Medicine)
Last reviewed: October 2026

I THE PARADIGM SHIFT (COMPLEMENTARY ROLE OF CT, MRI, AND ECHO)

- **Tri-Modality Synergy:** While **Transthoracic Echocardiography (TTE)** remains the universal first-line bedside tool for intracardiac anatomy and **Cardiac MRI (CMR)** is the radiation-free gold standard for ventricular volumes, ejection fraction, valvular regurgitant fractions (phase-contrast flow), and myocardial fibrosis, **Multidetector Cardiac Computed Tomography Angiography (MDCTA)** is the premier non-invasive modality for **extracardiac thoracic vasculature, coronary artery anomalies, tracheobronchial airways, and metallic stents/calcified conduits**.
- **Technical Edge of Modern CT:** Sub-millimeter **isotropic spatial resolution (0.4 to 0.6 mm)** and ultra-fast acquisition speeds (high-pitch dual-source spiral scanning or wide-detector volumetric scanning completing a whole-chest scan in a single heartbeat / less than 0.25 seconds) provide motion-free 3D reconstructions even at rapid neonatal heart rates.

II CORE PRE-OPERATIVE DIAGNOSTIC INDICATIONS (CLASS I)

- **Coronary Artery Anomalies (Modality of Choice — Class I):** MDCTA is the non-invasive reference standard for defining the ostial origin, proximal angulation, intramural segment, and termination of congenital coronary anomalies:
- **Anomalous Aortic Origin of a Coronary Artery (AAOCA):** Differentiates high-risk anatomical features—specifically an **anomalous Left Coronary Artery from the Right Sinus (ALCA)** or Right Coronary Artery from the Left Sinus (ARCA) taking a **malignant interarterial course** (between the aorta and pulmonary trunk) with a **slit-like ostium, acute take-off angle (less than 45 degrees), and intramural aortic wall course**—from a benign prepulmonic or retroaortic course.
- **Pre-Surgical TOF Planning:** Identifies an **anomalous LAD arising from the RCA (or large conal branch) crossing the Right Ventricular Outflow Tract (RVOT)** in Tetralogy of Fallot (present in 5 percent of cases), which precludes a standard transannular ventriculotomy incision.
- **ALCAPA and Coronary Fistulae:** Delineates **Anomalous Left Coronary Artery from the Pulmonary Artery (ALCAPA / Bland-White-Garland syndrome)** and complex **coronary cameral or arteriovenous fistulae**.
- **Aortic Arch Anomalies and Coarctation:** Provides 3D visualization of **Coarctation of the Aorta (CoA)**, **Interrupted Aortic Arch (IAA Types A, B, and C)**, **Truncus Arteriosus**, **Aortopulmonary Window**, and syndromic aortopathies (Marfan, Loeys-Dietz, Bicuspid Aortic Valve, and Turner syndrome).
- **Pulmonary Arterial Tree and MAPCAs:** Essential in **Pulmonary Atresia with VSD (TOF-PA)** to map the presence and confluence of central pulmonary arteries (calculating the **Nakata Index** and **McGoon Ratio**) and the exact aortic/branch origin, retroesophageal course, and lobar/segmental arborization of **Major Aortopulmonary Collateral Arteries (MAPCAs)** prior to surgical unifocalization.
- **Vascular Rings and Pulmonary Artery Slings (Simultaneous Vascular + Airway Assessment):** In infants presenting with biphasic stridor, wheezing, apnea, or dysphagia, a single CT acquisition simultaneously reconstructs **both the great vessels and the tracheobronchial lumen** (virtual bronchoscopy), diagnosing **Double Aortic Arch**, **Right Aortic Arch with Aberrant Left Subclavian Artery and Kommerell Diverticulum**, **Innominate Artery Compression**, and **Left Pulmonary Artery Sling** (associated with complete cartilaginous tracheal rings / “ring-sling complex”).
- **Anomalous Pulmonary and Systemic Venous Connections:** Gold standard for non-invasively mapping **Total Anomalous Pulmonary Venous Connection (TAPVC — Supracardiac, Cardiac, Infracardiac, and Mixed types)** and **Partial Anomalous Pulmonary Venous Connection (PAPVC / Scimitar Syndrome)**, precisely identifying vertical vein course and **pulmonary venous obstruction** (especially in infracardiac TAPVC draining into the portal/hepatic veins), as well as heterotaxy/isomerism systemic venous anomalies (interrupted IVC with azygos continuation, bilateral SVCs).

Clinical CHD Scenario	Preferred Modality (Cardiac CT vs. Cardiac MRI)	Key Diagnostic Advantage
Coronary Anomalies (AAOCA, ALCAPA, TOF crossing LAD)	**Cardiac CT (Class I Gold Standard)**	Superior sub-millimeter spatial resolution defines slit-like ostia and intramural course
Vascular Rings / Slings & Airway Compression	**Cardiac CT**	Simultaneously images great vessels **and tracheobronchial tree / lung parenchyma**
Neonatal Obstructed TAPVC & MAPCA Mapping	**Cardiac CT**	Sub-second scan in critically ill neonates **without general anesthesia**
Metallic Stents (CoA/PA), Calcified Conduits, Pacemakers	**Cardiac CT**	Free of the ferromagnetic susceptibility/drop-out artifacts that obscure MRI
RV Volumes/RVEF & PR Fraction in Repaired TOF	**Cardiac MRI (Class I Gold Standard)**	Radiation-free 3D cine volumetry + phase-contrast flow + myocardial LGE (CT used if non-MR conditional CIED)

III POST-OPERATIVE AND INTERVENTIONAL EVALUATION

- **Patency of Surgical Shunts, Conduits, and Baffles:** Superior to CMR for evaluating lumen patency and thrombosis of **Modified Blalock-Thomas-Taussig (mBTT) shunts**, **Sano conduits**, **Bidirectional Glenn** and **Fontan (TCPC) pathways** (using biphasic or dual-injection SVC/IVC contrast protocols to avoid unopacified streaming artifacts mimicking thrombus), and **Mustard/Senning intra-atrial baffles**.
- **Evaluation of Endovascular Stents and Conduit Calcification:** Modality of choice for assessing **in-stent intimal hyperplasia/restenosis, stent fracture, or pseudoaneurysm** in stented aortic coarctations, ductal/RVOT stents, or branch pulmonary artery stents (where stainless steel or cobalt-chromium stents cause severe black-blood signal drop-out on MRI), and for quantifying **circumferential conduit calcification** prior to redo sternotomy or balloon dilation.
- **Pre-Procedural Planning for Transcatheter Pulmonary Valve Replacement (TPVR):** MDCTA defines landing-zone dimensions of the RVOT/conduit for self-expanding (**Harmony, Alterra**) or balloon-expandable (**Melody, SAPIEN**) valves and assesses the **3D proximity of the proximal coronary arteries to the calcified RVOT conduit** to predict the risk of fatal **coronary artery compression** upon stent expansion, as well as enabling **3D printing** and computational modeling for complex double-outlet right ventricle (DORV) intracardiac baffling.

IV TECHNICAL ADVANTAGES, RADIATION SAFETY AND PATIENT SELECTION

- **Avoidance of General Anesthesia in Critically Ill Neonates:** Unlike a 45-to-60-minute CMR examination that requires general anesthesia and breath-holding in children under 6 to 7 years of age, modern **high-pitch Dual-Source CT ("Flash" mode)** or wide-detector CT acquires the entire thorax in **less than 0.25 to 0.5 seconds** during free breathing using a simple "feed-and-swaddle" technique or minimal oral/IV sedation.
- **Patients with Contraindications to MRI:** ECG-gated 4D cine Cardiac CT accurately quantifies **RV and LV volumes and ejection fraction** when CMR is contraindicated due to **non-MR-conditional epicardial/abandoned pacing leads, older pacemakers/ICDs, or ferromagnetic vascular coils**.
- **ALARA Radiation Dose Optimization:** Utilizing prospective ECG-triggering, high-pitch spiral scanning, low tube voltage (**70 to 80 kVp** in infants), and iterative reconstruction lowers pediatric cardiac CT effective radiation doses to **less than 0.5 to 1.0 mSv** (comparable to 10 to 20 chest radiographs).

V PRACTICAL CONSIDERATIONS IN THE INDIAN CLINICAL CONTEXT

- **Rapid Triage in Cyanotic Neonates and Malnourished Infants:** In Indian neonatal ICUs where critically ill neonates with **obstructed TAPVC** or **duct-dependent PA-VSD** carry prohibitive risks from prolonged general anesthesia, ultra-fast low-dose **CT Angiography** rapidly provides a definitive surgical roadmap without intubation.
- **Pragmatic Resource Allocation:** Because 256-slice and dual-source CT scanners are far more widely available across Indian Tier-1 and Tier-2 cities than dedicated congenital Cardiac MRI programs, MDCTA serves as the primary cross-sectional workhorse for pre-operative mapping of **MAPCAs, arch anomalies, and anomalous venous drainage**, while reserving specialized CMR slots for serial volumetric follow-up of repaired Tetralogy of Fallot and Fontan survivors.

KEY REFERENCES

- **Baumgartner H, De Backer J, Babu-Narayan SV, et al.:** 2020 ESC Guidelines for the management of adult congenital heart disease, *European Heart Journal*, 2020.
- **Han BK, Rigsby CK, Hlavacek A, et al.:** Computed Tomography Imaging in Patients with Congenital Heart Disease Part I & Part II: An Expert Consensus Document of the Society of Cardiovascular Computed Tomography (SCCT), *Journal of Cardiovascular Computed Tomography*, 2015.
- **Sachdeva R, Valente AM, Armstrong AK, et al.:** ACC/AHA/ASE/HRS/ISACHD/SCAI/SCCT/SCMR/SOPE 2020 Appropriate Use Criteria for Multimodality Imaging During the Follow-Up Care of Patients With Congenital Heart Disease, *Journal of the American College of Cardiology*, 2020.