SINUS NODE FUNCTION ASSESSMENT: CLINICAL AND ELECTROPHYSIOLOGICAL PROTOCOLS

Prepared by **Dr. Sidhartha Mullapudi**, DrNB (Cardiology), MD (General Medicine)
Last reviewed: October 2026

I NON-INVASIVE EVALUATION (THE CLINICAL BASELINE)

- **Principle:** Invasive testing is rarely the first step. Symptom-rhythm correlation is the absolute gold standard for diagnosing Sinus Node Dysfunction (**SND**).
- **Holter Monitoring:** Essential to document sinus pauses (greater than 3 seconds), severe inappropriate bradycardia, or Tachy-Brady Syndrome.
- **Treadmill Exercise Testing:** Used specifically to diagnose **Chronotropic Incompetence**, defined as the inability to reach 85 percent of the age-predicted maximum heart rate during peak exercise.
- **Intrinsic Heart Rate (IHR):** Assessed via pharmacological autonomic blockade to determine if the sinus node is intrinsically diseased or just under excessive vagal tone.
  - **Protocol:** Administer intravenous Propranolol (0.2 mg/kg) followed by Atropine (0.04 mg/kg).
  - **Calculation:** Normal **IHR** = 118.1 – (0.57 × age). An **IHR** significantly lower than calculated proves intrinsic structural disease of the Sinus Node.

II INVASIVE ELECTROPHYSIOLOGY STUDY (EPS): THE CORE EXAM METRICS

- **When to Perform:** When non-invasive tests are inconclusive in a highly symptomatic patient, an **EPS** is performed to directly measure sinus node automaticity and conduction.

A Sinus Node Recovery Time (SNRT)

- **Physiology:** Rapid atrial pacing suppresses the Sinus Node (overdrive suppression). When pacing is abruptly stopped, the time it takes for the Sinus Node to “wake up” and fire the first spontaneous P wave is the **SNRT**.
- **Protocol:** Pace the High Right Atrium (**HRA**) at a fixed cycle length (e.g., 600 ms) for 30 to 60 seconds, then suddenly terminate.
- **Normal Value:** Less than 1,500 ms.

B Corrected Sinus Node Recovery Time (cSNRT)

- **Physiology:** The raw **SNRT** is heavily influenced by the patient’s baseline resting heart rate. The **cSNRT** mathematically removes the baseline rate to provide the true recovery time.
- **Calculation: SNRT** minus basic Sinus Cycle Length (**SCL**).
- **Normal Value:** Less than 525 ms. A prolonged **cSNRT** is highly specific for Sick Sinus Syndrome.

C Sinoatrial Conduction Time (SACT)

- **Physiology:** The time it takes for the electrical impulse generated within the Sinus Node to exit the node and depolarize the surrounding right atrial tissue.
- **Narula Method:** Continuous rapid atrial pacing just slightly faster than the resting sinus rate.
- **Strauss Method:** Delivering premature atrial extrastimuli during a stable sinus rhythm.
- **Normal Value:** Less than 125 ms. Prolonged **SACT** indicates Sinoatrial Exit Block.

III PATHOPHYSIOLOGY AND CLINICAL CORRELATION

- **Sick Sinus Syndrome (SSS):** A degenerative, fibrotic disease of the Sinus Node and surrounding atrial myocardium. Often presents with alternating periods of profound bradycardia and rapid atrial arrhythmias (Atrial Fibrillation).
- **Sinoatrial Exit Block:** The node generates an impulse normally, but fibrosis prevents the impulse from reaching the atrium.
  - **Type One (Wenckebach):** Progressive shortening of the P-to-P interval until a P wave drops.
  - **Type Two:** Sudden dropped P wave. The pause is an exact multiple of the baseline P-to-P interval.

IV THE CLINICAL REALITY

- **Pacing Decisions:** In routine practice, from district clinics to advanced tertiary centers, the decision to implant a permanent pacemaker for **SND** is almost never based solely on an abnormal **cSNRT** or **SACT** value. The Class One indication strictly requires documented symptomatic bradycardia or symptomatic chronotropic incompetence. If the patient is asymptomatic, even a markedly prolonged **cSNRT** during an **EPS** does not justify pacing.